

The content on the UpToDate website is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your own physician or other qualified health care professional regarding any medical questions or conditions. The use of UpToDate content is governed by the [UpToDate Terms of Use](#). ©2016 UpToDate, Inc. All rights reserved.

Patient education: Hypothyroidism (underactive thyroid) (Beyond the Basics)

Author: [Douglas S Ross, MD](#)

Section Editor: [David S Cooper, MD](#)

Deputy Editor: [Jean E Mulder, MD](#)

All topics are updated as new evidence becomes available and our [peer review process](#) is complete.

Literature review current through: Aug 2016. | **This topic last updated:** Mar 24, 2016.

HYPOTHYROIDISM OVERVIEW — Hypothyroidism is a condition in which the thyroid gland does not produce enough thyroid hormone. It is the most common thyroid disorder.

This topic discusses **hypothyroidism**. **hyperthyroidism** is discussed separately. (See "[Patient education: Hyperthyroidism \(overactive thyroid\) \(Beyond the Basics\)](#)".)

WHAT IS THE THYROID? — The thyroid is a butterfly-shaped gland in the middle of the neck, located below the larynx (voice box) and above the clavicles (collarbones) ([figure 1](#)). The thyroid produce two hormones, triiodothyronine (T3) and thyroxine (T4), which regulate how the body uses and stores energy (also known as the body's metabolism).

Thyroid function is controlled by a gland in the brain, known as the pituitary ([figure 2](#)). The pituitary produces thyroid-stimulating hormone (TSH), which stimulates the thyroid to produce T3 and T4.

HYPOTHYROIDISM CAUSES — In about 95 percent of cases, hypothyroidism is due to a problem in the thyroid gland itself and is called primary hypothyroidism. However, certain medications and diseases can also decrease thyroid function. As an example, **hypothyroidism** can also develop after medical treatments for **hyperthyroidism**, such as thyroidectomy (surgical removal of the thyroid) or radioactive iodine treatment (to destroy thyroid tissue). In some cases, hypothyroidism is a result of decreased production of thyroid-stimulating hormone (TSH) by the pituitary gland (called secondary hypothyroidism). (See "[Patient education: Hyperthyroidism \(overactive thyroid\) \(Beyond the Basics\)](#)".)

Thyroid problems are more common in women, increase with age, and are more common in whites and Mexican Americans than in blacks.

HYPOTHYROIDISM SYMPTOMS — The symptoms of hypothyroidism vary widely; some people have no symptoms while others have dramatic symptoms or, rarely, life-threatening symptoms. The symptoms of hypothyroidism are notorious for being nonspecific and for mimicking many of the normal changes of aging. Usually, symptoms are milder when hypothyroidism develops gradually. Symptoms, when caused by hypothyroidism, generally are related to the degree of hypothyroidism. Many patients with mild hypothyroidism are identified on screening tests for potential hypothyroid symptoms, but have few or no symptoms that ultimately are attributed to hypothyroidism or respond to treatment of hypothyroidism. In contrast, patients with moderate to severe hypothyroidism are usually symptomatic and improve significantly with thyroid hormone replacement.

The following list of symptoms are those that may be present prior to treatment; hypothyroid patients on appropriate treatment should no longer be symptomatic. If symptoms persist, they likely have causes other than hypothyroidism.

General symptoms — Thyroid hormone normally stimulates the metabolism, and most of the symptoms of hypothyroidism reflect slowing of metabolic processes. General symptoms may include fatigue, sluggishness, weight gain, and intolerance of cold temperatures.

Skin — Hypothyroidism can decrease sweating. The skin may become dry and thick. The hair may become coarse or thin, eyebrows may disappear, and nails may become brittle.

Eyes — Hypothyroidism can lead to mild swelling around the eyes. People who develop hypothyroidism after treatment for Graves' disease may retain some of the eye symptoms of Graves' disease, including protrusion of the eyes, the appearance of staring, and impaired movement of the eyes. (See ["Patient education: Hyperthyroidism \(overactive thyroid\) \(Beyond the Basics\)"](#).)

Cardiovascular system — Hypothyroidism slows the heart rate and weakens the heart's contractions, decreasing its overall function. Related symptoms may include fatigue and shortness of breath with exercise. These symptoms may be more severe in people who also have heart disease. In addition, hypothyroidism can cause mild high blood pressure and raise blood levels of cholesterol.

Respiratory system — Hypothyroidism weakens the respiratory muscles and decreases lung function. Symptoms can include fatigue, shortness of breath with exercise, and decreased ability to exercise. Hypothyroidism can also lead to swelling of the tongue, hoarse voice, and sleep apnea. Sleep apnea is a condition in which there is intermittent blockage of the airway while sleeping, causing fitful sleep and daytime sleepiness. (See ["Patient education: Sleep apnea in adults \(Beyond the Basics\)"](#).)

Gastrointestinal system — Hypothyroidism slows the actions of the digestive tract, causing constipation. Rarely, the digestive tract may stop moving entirely. (See ["Patient education: Constipation in adults \(Beyond the Basics\)"](#).)

Reproductive system — Women with hypothyroidism often have menstrual cycle irregularities, ranging from absent or infrequent periods to very frequent and heavy periods. The menstrual irregularities can make it difficult to become pregnant, and pregnant women with hypothyroidism have an increased risk for miscarriage during early pregnancy. Treatment of hypothyroidism can decrease these risks. (See ["Patient education: Absent or irregular periods \(Beyond the Basics\)"](#) and ["Patient education: Heavy or prolonged menstrual bleeding \(menorrhagia\) \(Beyond the Basics\)"](#).)

Myxedema coma — In people with severe hypothyroidism, trauma, infection, exposure to the cold, and certain medications can rarely trigger a life-threatening condition called myxedema coma, which causes a loss of consciousness and hypothermia (low body temperature).

HYPOTHYROIDISM DIAGNOSIS — In the past, hypothyroidism was not diagnosed until symptoms had been present for a long time. However, simple blood tests can now detect hypothyroidism at an early stage. A person may be tested for hypothyroidism if there are signs and symptoms, such as those discussed above, or as a screening test.

Blood tests — Blood tests can confirm the diagnosis and pinpoint the underlying cause of the thyroid hormone deficiency. The most common blood test for hypothyroidism is thyroid-stimulating hormone (TSH). TSH is the most sensitive test because it can be elevated even with small decreases in thyroid function. Thyroxine (T4), the main product of the thyroid gland, may also be measured to confirm and assess the degree of hypothyroidism.

Routine screening — All newborn babies in the United States are routinely screened for thyroid hormone deficiency. It is not clear if all adults should be tested for thyroid disease [1].

HYPOTHYROIDISM TREATMENT — The goal of treatment for hypothyroidism is to return blood levels of thyroid-stimulating hormone (TSH) and thyroxine (T4) to the normal range and to alleviate symptoms.

Medication — The treatment for hypothyroidism is thyroid hormone replacement therapy. This is usually given as an oral form of T4 (levothyroxine). T4 should be taken once per day on an empty stomach (ideally one hour before eating or two hours after; most patients take their hormone as soon as they wake in the morning, and delay eating breakfast as long as practical before leaving for work or school). Generic (levothyroxine) and brand-name (Synthroid, Levoxyl, Levothroid, Unithyroid, Tirosint) formulations are equally effective. However, it is preferable to stay on the same manufacturer of T4 rather than switching between brand name and/or generic formulations.

If a switch is necessary and the patient feels that their levels may be off, a blood test can be done six weeks later to determine if the dose needs to be adjusted. Color-coded tablets can help with dose adjustments.

Some clinicians prescribe another form of thyroid hormone, T3 (liothyronine), in combination with T4. However, since T4 is converted into triiodothyronine (T3) in other organs, the majority of studies have not shown an advantage of combination T3 and T4 therapy over T4 alone.

In most cases, symptoms of hypothyroidism begin to improve within two weeks of starting thyroid replacement therapy. However, people with more severe symptoms may require several months of treatment before they fully recover.

Duration and dose — A healthcare provider will prescribe an initial dose of T4 and then retest the blood level of TSH after six weeks. The T4 dose can be adjusted at that time, depending upon these results. This process may be repeated several times before hormone levels become normal. After the optimal dose is identified, a provider may recommend monitoring blood tests once yearly, or more often as needed. Most people with hypothyroidism require lifelong treatment, although the dose of T4 may need to be adjusted over time.

Never increase or decrease the T4 dose without first consulting a healthcare provider. Overreplacement of T4 can cause mild hyperthyroidism, with the associated dangers of atrial fibrillation (irregular heart beat) and, possibly, accelerated bone loss (osteoporosis).

Dose changes — Changes in the T4 dose are based upon the person's TSH and T4 level. The dose may need to be increased if thyroid disease worsens, during pregnancy, if gastrointestinal conditions impair T4 absorption, or if the person gains weight. A high fiber diet, calcium- or aluminum-containing antacids, and iron tablets can interfere with the absorption of T4 and should be taken at a different time of day.

The dose may need to be decreased as the person gets older, after childbirth, or if the person loses weight.

Monitoring — Individual T4 doses can vary widely and depend upon a variety of factors, including the underlying cause of hypothyroidism. People with certain conditions require more frequent monitoring.

Advanced age and heart disease — Thyroid hormone makes the heart work a bit harder. Therefore, a clinician may opt for more conservative T4 treatment in older adults and in people with coronary artery disease.

Pregnancy — Women often need higher doses of T4 during pregnancy. Testing is usually recommended every four weeks, beginning after conception, until levels are stable, then once each trimester. After delivery, the woman's dose of T4 will need to be adjusted again, usually returning to the pre-pregnancy dose.

Surgery — Hypothyroidism can increase the risk of certain surgery-related complications; bowel function may be slow to recover and infection may be overlooked if there is no fever. If pre-operative blood tests reveal low thyroid hormone levels, non-emergency surgery is usually postponed until treatment has returned T4 levels to normal.

Hypothyroidism without symptoms — In some cases, hypothyroidism is extremely mild or causes no obvious symptoms (called subclinical hypothyroidism). The decision to treat subclinical hypothyroidism with T4 is controversial. Many experts treat patients with subclinical hypothyroidism if their TSH is >7 to 10 mU/L to prevent the development of hypothyroidism and associated symptoms. Treatment is also recommended for people who have a goiter or nonspecific symptoms of hypothyroidism, such as fatigue, constipation, or depression.

WHERE TO GET MORE INFORMATION — Your healthcare provider is the best source of information for questions and concerns related to your medical problem.

This article will be updated as needed on our web site (www.uptodate.com/patients). Related topics for patients, as well as selected articles written for healthcare professionals, are also available. Some of the most relevant are listed below.

Patient level information — UpToDate offers two types of patient education materials.

The Basics — The Basics patient education pieces answer the four or five key questions a patient might have about a given condition. These articles are best for patients who want a general overview and who prefer short, easy-to-read materials.

[Patient education: Hypothyroidism \(underactive thyroid\) \(The Basics\)](#)

[Patient education: Hemochromatosis \(The Basics\)](#)

[Patient education: Thyroid nodules \(The Basics\)](#)

[Patient education: Congenital hypothyroidism \(The Basics\)](#)

[Patient education: Hypoparathyroidism \(The Basics\)](#)

[Patient education: Panhypopituitarism \(The Basics\)](#)

[Patient education: Thyroiditis after pregnancy \(The Basics\)](#)

[Patient education: Thyroiditis \(The Basics\)](#)

Beyond the Basics — Beyond the Basics patient education pieces are longer, more sophisticated, and more detailed. These articles are best for patients who want in-depth information and are comfortable with some medical jargon.

[Patient education: Hyperthyroidism \(overactive thyroid\) \(Beyond the Basics\)](#)

[Patient education: Sleep apnea in adults \(Beyond the Basics\)](#)

[Patient education: Constipation in adults \(Beyond the Basics\)](#)

[Patient education: Absent or irregular periods \(Beyond the Basics\)](#)

[Patient education: Heavy or prolonged menstrual bleeding \(menorrhagia\) \(Beyond the Basics\)](#)

Professional level information — Professional level articles are designed to keep doctors and other health professionals up-to-date on the latest medical findings. These articles are thorough, long, and complex, and they contain multiple references to the research on which they are based. Professional level articles are best for people who are comfortable with a lot of medical terminology and who want to read the same materials their doctors are reading.

[Acquired hypothyroidism in childhood and adolescence](#)

[Cardiovascular effects of hypothyroidism](#)

[Central hypothyroidism](#)

[Clinical features and detection of congenital hypothyroidism](#)

[Clinical manifestations of hypothyroidism](#)

[Diagnosis of and screening for hypothyroidism in nonpregnant adults](#)

[Disorders that cause hypothyroidism](#)

[Laboratory assessment of thyroid function](#)

[Myxedema coma](#)

[Neurologic manifestations of hypothyroidism](#)

[Overview of thyroid disease in pregnancy](#)

[Subclinical hypothyroidism in nonpregnant adults](#)

[Treatment and prognosis of congenital hypothyroidism](#)

[Treatment of hypothyroidism](#)

[Endocrinopathies in the childhood cancer survivor](#)

The following organizations also provide reliable health information.

- National Library of Medicine

(www.nlm.nih.gov/medlineplus/healthtopics.html)

- The American Thyroid Association

(www.thyroid.org)

- Hormone Health Network

(www.hormone.org/diseases-and-conditions/thyroid/hypothyroidism)

Patient support — There are a number of online forums where patients can find information and support from other people with similar conditions.

- [About.com](http://thyroid.about.com/forum) Thyroid Conditions Forum

(<http://thyroid.about.com/forum>)


[1.2]

Use of UpToDate is subject to the [Subscription and License Agreement](#).

Topic 2179 Version 11.0

GRAPHICS

Thyroid and parathyroid glands




The thyroid is a butterfly-shaped gland in the middle of the neck. It sits just below the larynx (voice box). The thyroid makes two hormones, called T3 and T4, which control how the body uses and stores energy. The parathyroid glands are four small glands behind the thyroid. They make a hormone called parathyroid hormone, which helps control the amount of calcium in the blood.

T3: triiodothyronine; T4: thyroxine.

Graphic 66834 Version 8.0

Endocrine glands



This figure shows the location of the adrenal glands, hypothalamus, and pituitary.

Graphic 67279 Version 4.0

Contributor Disclosures

Douglas S Ross, MD Consultant/Advisory Board: Eisai [Thyroid cancer (Lenvatinib)]; Novo Nordisk [Thyroid cancer]; Medullary Thyroid Cancer Registry Consortium [Thyroid cancer]. **David S Cooper, MD** Nothing to disclose **Jean E Mulder, MD** Nothing to disclose

Contributor disclosures are reviewed for conflicts of interest by the editorial group. When found, these are addressed by vetting through a multi-level review process, and through requirements for references to be provided to support the content. Appropriately referenced content is required of all authors and must conform to UpToDate standards of evidence.

[Conflict of interest policy](#)

[Close](#)