



Patient education: Flu (The Basics)

[Written by the doctors and editors at UpToDate](#)

What is the flu? — The flu is an infection that can cause fever, cough, body aches, and other symptoms. There are different forms of the flu. The most common is the "seasonal" flu. There is also the "swine" flu, which caused a worldwide outbreak ("pandemic") in 2009 and 2010, and the bird flu. Bird flu (also known as "avian flu") is a severe form of the flu that is caused by a type of flu virus that first infected birds.

All forms of the flu are caused by viruses. The medical term for the flu is "influenza."

What are the most common flu symptoms? — All forms of the flu can cause:

- Fever (temperature higher than 100°F or 37.8°C)
- Cough
- Headache or body aches

Less common symptoms are sore throat and a runny nose.

Is the flu dangerous? — It can be. Most people get over the flu on their own, without any lasting problems. But some people need to go to the hospital because of the flu. And some people even die from it. This is because the flu can cause a serious lung infection called pneumonia. That's why it's important to keep from getting the flu in the first place.

Is there a test for flu? — Yes. There are tests for the flu. In most cases, your doctor can tell if you have the flu by your symptoms. But in some cases – for example, if you are at risk for having other problems caused by the flu – your doctor might do a test for flu.

How can I protect myself from the flu? — You can:

- Wash your hands often with soap and water, or use alcohol hand rubs
- Stay away from people you know are sick
- Get the flu vaccine every year – There are a few different kinds of flu vaccine. Your doctor can help you decide which vaccine is best for you.

What should I do if I get the flu? — If you think you have the flu, stay home, rest, and drink plenty of fluids. You can also take [acetaminophen](#) (sample brand name: Tylenol) to relieve fever and aches.

Do not give [aspirin](#) or medicines that contain aspirin to children younger than 18. In children, aspirin can cause a serious problem called Reye syndrome.

Most people with the flu get better on their own within 1 to 2 weeks. But you should call your doctor or nurse if you:

- Have trouble breathing or are short of breath
- Feel pain or pressure in your chest or belly
- Get suddenly dizzy
- Feel confused
- Have severe vomiting

Take your child to the doctor if he or she:

- Starts breathing fast or has trouble breathing
- Starts to turn blue or purple
- Is not drinking enough fluids
- Will not wake up or will not interact with you
- Is so unhappy that he or she does not want to be held
- Gets better from the flu but then gets sick again with a fever or cough
- Has a fever with a rash

If you decide to go to a walk-in clinic or a hospital because of the flu, tell someone right away why you are there. The staff might ask you to wear a mask or to wait someplace where you are less likely to spread your infection.

Whether or not you see a doctor or nurse, you should stay home while you are sick with the flu. Do not go to work or school until your fever has been gone for at least 24 hours, without taking fever-reducing medicine, such as [acetaminophen](#). If you work in a healthcare setting taking care of patients, you might need to stay home longer if you are still coughing. Also, always cover your mouth and nose with the inside of your elbow when you cough or sneeze.

Can the flu be treated? — Yes, people with the flu can get medicines called antiviral medicines. These medicines can help people avoid some of the problems caused by the flu. Not every person with the flu needs an antiviral medicine, but some people do. Your doctor or nurse will decide if you need an antiviral medicine. Antibiotics **do not work** on the flu.

What if I am pregnant? — The flu can be very dangerous for pregnant women. If you are pregnant, it is very important that you get the flu vaccine. You should also avoid taking care of anyone who has the flu.

If you are pregnant, call your doctor or nurse right away if:

- You might have been near someone with the flu.
- You think you might be coming down with the flu. In pregnant women, the symptoms of the flu can get worse very quickly. The flu can even cause trouble breathing or lead to death of the woman or her baby. That is why it is so important that you talk to doctor or nurse as soon as you notice any of the flu symptoms listed above. You will need an antiviral medicine if you are pregnant and have the flu.

More on this topic

[Patient education: Flu vaccine \(The Basics\)](#)

[Patient education: Cough, runny nose, and the common cold \(The Basics\)](#)

[Patient education: Fever in children \(The Basics\)](#)

[Patient education: Vaccines \(The Basics\)](#)

[Patient education: Sore throat in adults \(The Basics\)](#)

[Patient education: Giving your child over-the-counter medicines \(The Basics\)](#)

[Patient education: What you should know about antibiotics \(The Basics\)](#)

[Patient education: When to worry about a fever in adults \(The Basics\)](#)

[Patient education: Bird flu \(avian influenza\) \(The Basics\)](#)

[Patient education: Influenza symptoms and treatment \(Beyond the Basics\)](#)

[Patient education: Influenza prevention \(Beyond the Basics\)](#)

All topics are updated as new evidence becomes available and our [peer review process](#) is complete.

This topic retrieved from UpToDate on: Dec 22, 2017.

The content on the UpToDate website is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your own physician or other qualified health care professional regarding any medical questions or conditions. The use of UpToDate content is governed by the [UpToDate Terms of Use](#). ©2017 UpToDate, Inc. All rights reserved.

Topic 15369 Version 11.0